



ELSEVIER

Review

Mental distress through the prism of predictive processing theory

Sander Van de Cruys^{1,2} and Pieter Van Dessel³

Summary

We review the predictive processing theory's take on goals and affect, to shed new light on mental distress and how it develops into psychopathology such as in affective and motivational disorders. This analysis recovers many of the classical factors known to be important in those disorders, like uncertainty and control, but integrates them in a mechanistic model of adaptive and maladaptive cognition and behavior. We derive implications for treatment that have so far remained underexposed in existing predictive processing accounts of mental disorder, specifically with regard to the model-dependent construction of value, the importance of model validation (evidence), and the introduction and learning of new, adaptive beliefs that relieve suffering.

Addresses

¹Laboratory of Experimental Psychology, KU Leuven, Belgium²Antwerp Social Lab, University of Antwerp, Belgium³Department of Experimental-Clinical and Health Psychology, Ghent University, Belgium

Corresponding authors: Pieter Van Dessel (Pieter.VanDessel@UGent.be); Sander Van De Cruys (Sander.Vandecruys@kuleuven.be)

Current Opinion in Psychology 2021, **41**:107–112

This review comes from a themed issue on **Psychopathology**

Edited by **Peter de Jong** and **Yannick Boddez**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 19 July 2021

<https://doi.org/10.1016/j.copsyc.2021.07.006>

2352-250X/© 2021 Elsevier Ltd. All rights reserved.

Keywords

Predictive processing, Mental distress, Psychopathology, Emotion, Depression, Anxiety, Active inference, Addiction, Learning, Psychotherapy, Computational psychiatry.

Introduction

In recent decades, the influential theory of predictive processing (PP) (also called Bayesian brain or active inference) [1–3] has increasingly gained traction. It states that organisms do not merely respond to stimuli but proactively model and predict sensory inputs and the opportunities and challenges they represent. This view implements pioneering ideas of Jerome Bruner [4] and George Kelly [5] that describe mental processing in

terms of the formation and testing of hypotheses. As true proto-scientists, we make predictions about the world and meet reality only in our failures [6], that is, when sensory input deviates from our predicted or constructed versions. Such deviations are called *prediction errors* in PP lingo. Because we perceive through our own constructs (in the form of predictions), perception is often described as ‘controlled hallucination’, which means that we de facto live in a constructed reality [7], kept in check by prediction errors.

The computational details of PP are beyond the current scope [8–10], but the central thesis is that the mental system realizes all its functions through prediction error minimization. Organisms embody an evolutionarily ‘discovered’ set of expected interoceptive (homeostatic) states that are required for continued existence but that they can only realize via action in their environment (e.g. foraging for food to increase blood glucose levels). This requires them to build a generative model about the structure of their environment and how to ‘generate’ their expected states through action. This model allows them to not only react on homeostatic disturbances, but also predict and so preempt those. It is shaped by prediction errors, which can be minimized either by acting to bring predicted events in line with the current model (cf. assimilation), or by perception and learning, that is, by changing our models and inferences so they correspond better with incoming evidence (cf. accommodation). In this way, the organism maximizes the evidence for the model it embodies, often called *self-evidencing* [11]. Importantly, our sense of self can be seen as just another inferred cause in the generative model, as the best explanation for the regularities in multimodal perceptual consequences created by one's own actions, and as an efficient prediction of (regularities in) one's future behavior, emotions, and thoughts (e.g. the narrative and social self). From this perspective, we construe and infer not only what is there in the world, but also who we are, what we can do, and what we want (i.e. our goals), as also described in self-perception theories [12].

The PP framework hence entails that a person's behavior is canalized by the ways in which they construe and anticipate events and their own reactions to these (constructed) events [5]. Evidently, this holds for mental distress and disorder too. It is clear that people can end up with sometimes drastically diverging

constructs (or beliefs) and mental models, first because of the vagaries of the environment that shape the models (i.e. their phylogenetic and ontogenetic learning history), second, because the updating of constructs using prediction errors is a fallible job that requires weighing evidence based on different types of uncertainty [13], and third, because the very gathering of evidence is steered (biased) by the current constructs. Because of its unique capacity to account for the variety and idiosyncrasy of models, we will argue that the PP framework is well placed to help us understand mental distress and, hopefully, its treatment.

Goals in predictive processing

The overarching principle of prediction error minimization means that the mental system is geared toward *model validation*, maximizing the evidence for one's model or, equivalently, reducing uncertainty relative to this model. At first glance, this conflicts with established reward-based or goal-directed theories [14] that see reward maximization or goal realization as the core organizing principle for the mind. But, as pointed out above, both biological needs and more personal goals are folded into the model. Goals are expected states (desired outcomes) that can be attained ('generated') by our own actions. Some (core) goals are defined at a higher level (longer temporal horizon) and are persistent against obstacles, that is, these expectations will *not* be readily updated by momentary prediction errors. Evidently, this resistance is crucial for homeostatic expected states, where 'updates' (e.g. lowering glucose expectation) would lead to loss of organismic integrity. However, humans construe a myriad of other goals and values in (putative) support of those interoceptive expectations. We infer our goals and values from the history of our choices and actions (e.g. approach and avoidance) [15], as well as from what people like us (parents, peers, and mentors) do and want. What makes these expectations resistant to updates is the *precision* they have accumulated through experience. Precision is the technical term in PP for the estimate of expected uncertainty associated with expected sensory states, in this particular case the expected outcomes of actions. In the literature on goal-directed behavior [16], this expected uncertainty about mapping between actions and outcomes is known as 'feasibility' or control (lability).

Given that any behavior is driven by expected (interoceptive or exteroceptive) states, habits in the sense of automatic stimulus-response mappings have no place in PP. Instead, habit-like behavior can be accounted for as actions that are based on strong stimulus-stimulus predictions (one sensory input predicts another proprioceptive and exteroceptive input) maintained on hierarchically lower levels. Although these actions were originally driven by higher level goals, through experience, the lower level expectations involved (sub-goals) accumulated enough precision (reduced uncertainty) so

the behavior is now sustained by 'shallow models' [17,18]. In contrast, behavior commonly referred to as 'goal-directed behavior' depends on higher level goals and involves lower level expectations that are contextualized by hierarchically deep models. These higher level goals are only activated when there is uncertainty about actions and whether those lead to desired outcomes. For instance, when sensory input activates different alternative actions, deliberation will involve weighing whether one already has reliable (precise) actions to attain desired outcomes, or whether one needs to gather more information about the environment (including the consequences of one's actions) through so-called epistemic actions. This deliberation may remain implicit, but will include counterfactual beliefs: "What if I do this, instead of that, how well does that reduce my expected prediction errors (uncertainty)?" Hence, whether some behavior is goal-directed or habit-like, is a matter of degree, dependent on how shallow or (counterfactually) 'deep' the model is that led to this behavior.

This conceptualization of goals casts a new light on clinically relevant cases where verbally reported goals conflict with (possibly maladaptive) actual behavior that is more habit-like but possibly still driven by lower level 'goals' or desired outcomes [18]. Indeed, here we essentially have two different models within one agent applied to the same behavior: The model that supports the behavior of self-reporting our explicit, conscious goals, and the model that supports the habit-like behavior. For example, in addiction, a particular 'eliciting' context would evoke high precision beliefs about interoceptive expected consequences of substance use (the habit-like behavior). But such beliefs are distinct from the (more explicit) beliefs that underlie statements about this behavior and its consequences in a different context (e.g. during a therapy session). Note that the terms 'beliefs' and models are used in the Bayesian sense here, as implicit, inferred networks of hidden causes (technically: probability distributions over parameter values) or constructs that explain input and sustain behaviors [19]. Part of the work in clinical therapy, as psychotherapists can attest, is therefore to articulate these constructs underlying habit-like (pathological) behavior, not so much to be able to directly confront them with the clients' conscious goals, but to confront them with inconsistent evidence directed toward their own logic. This will boost the expected uncertainty of unwanted beliefs, enabling new learning and epistemic actions to establish constructs that support effective behavior in line with one's explicit goals. While this view confirms received views of cognitive behavioral therapy, it also acknowledges the problem of fragmented models: The fact that some (parts of) models that support problematic behavior are so 'inferentially isolated' that the task of un/relearning models cannot be undertaken right away, but should be preceded by a reconstruction of underlying beliefs and

desires. This quest to understand the subject's web of constructs (their generative model) from within is akin to an anthropologist's task [20], but it is required to target the right 'wrong' beliefs in therapy.

Mental distress in predictive processing

The PP framework has already been applied to several psychopathologies, such as autism spectrum disorder [21], schizophrenia [22], and obsessive-compulsive disorder [23], usually by characterizing these conditions as stemming from a general deficiency in prediction updating (technically: precision weighting). However, to explain mental distress in affective and motivational disorders with their strong link to life events, we first need to look at the way PP understands emotions.

Dominant views of affect cast it as a continuous "neurophysiological barometer of the individual's relationship to an environment at a given point in time" [24]. Similarly, Frijda [25] noted: "pleasure is the positive outcome of constantly monitoring one's functioning." Translated to PP, this kind of monitoring of one's own performance in predicting internal and external disturbances, needs to be an operation on prediction errors. However, given that prediction error minimization is really all the brain does (when perceiving, acting, and thinking), any momentary prediction error cannot be very informative or 'affective'. Instead, meta-expectations about the temporal average of prediction errors and how that changes—whether uncertainty increases or decreases relative to one's models—may fulfill this meta-monitoring function that we can associate with emotional valence [26–28]. This aligns with the well-known role of increasing uncertainty in negative affect such as anxiety [29], and of decreasing uncertainty in pleasure, such as aesthetic appreciation [30]. Specifically, given that, as described in the previous section, we absorb goals into the agent's model, decreasing/increasing uncertainties relative to that model are precisely informative about our progress/regress relative to our goals and the aptness of our models to deal with our surroundings.

This sets the stage for an understanding of distress as rooted in persistent breaches against predictive models, especially concerning high-level, high precision expected states (goals), because those predictions structure major parts of one's behavior, and are the basis for interpretation of much of the perceptual inputs the person encounters (past, present, and future). Persistent and increasing prediction errors signal the subversion of a person's constructed identity: the regularities from the world that they have absorbed but that no longer hold. It is here that we situate normal and pathological distress, be it grief, depression, or anxiety.

Usually, prediction errors (even with regard to core goals) can be neutralized by changing one's auxiliary assumptions, and so keeping the core of a person's models intact

(sometimes called cognitive immunization). That is just run-of-the-mill model updating which will take place first for lower level, less precise predictions, rather than for higher level, precise predictions. However, persistent model disconfirmations can incite a number of different responses to reduce uncertainty (increase model evidence). A first approach might be to tighten up on one's constructs, and insist on their validity. This is often expressed as a more selective data gathering to file off the sharp edges of disconfirmations, in other words: avoiding particular (e.g. social) situations and retreat to more reliably predictable environments, as we often see in the rituals of anxious people. Another outcome could be hostility, to be seen as extreme efforts to force the circumstances to conform to one's constructs again. Whether avoidant action or aggressive action is taken may depend on perceived control, a well-known factor in emotion and psychopathology, which, in PP, can be conceptualized as uncertainty about (our prediction of) action consequences [31].

Crucially, persistent prediction errors in reaching desired action outcomes can lead to model updates that increase the expected uncertainty or, simply put, the perceived powerlessness of actions. The generalized loss of confidence (certainty) that one can attain the expected consequences of one's own actions (cf. learned helplessness) can then lead to withdrawal and other depressive symptoms. This resonates with the fact that unpredictable, unresponsive environments, especially early on in development, are a key risk factor for (depressive) psychopathology [32].

The 'system-wide' generalization of loss of confidence in the efficacy of all actions (as in depression) might seem unwarranted but it may actually be internally rational, given one's models and the experiences that shaped them. Usually, the reverse generalization, a certain generalized overconfidence (known as the optimism bias or the overestimation of control) keeps us motivated and is often seen as a mark of good mental health [33]. However, a generalized *loss* of confidence in actions shifts one's mental model from trying to fit the world to their mental model (by actions) to fitting the mental model to the world and may therefore be a successful way to more easily achieve prediction error minimization. This might relate to depressive rumination (searching for better models without seeking out new evidence) as well as depressive realism, the finding that people with depression often have a more accurate perception of their capacities and control (i.e. a more accurate model of the present).

Finally, another effective strategy to deal with persistent prediction errors is to form predictions on a more global, less granular level [34]. Simply stated, categorizing something as merely 'fruit' or 'vegetable', rather than at a more specific level, allows one to be correct more

often. Applied to psychopathologies such as anxiety disorder, over-general threat-related priors (e.g. ‘social situations are threatening’) may start to dominate thought and behavior because they ‘worked’ (i.e. they were effective in reducing errors) and have therefore accumulated great (perceived) precision, hence becoming less sensitive to new contradicting evidence [35]. In addition, by putting all social situations under one umbrella, the lack of confidence associated with an action in one particular social situation easily propagates to lack of confidence in the ability of actions for social situations across the board. The one action that remains is avoidance, which of course prevents the confrontation with new evidence altogether.

Interestingly, none of these mechanisms to cope with persistent prediction errors can be labeled as pathological per se (they are part of the conspiracy theorist’s mind, and even the ‘normal’ mind). They are valid, adaptive ways of curbing prediction errors, not specific to psychopathologies. Hence our focus on distress in general. However, the salience of prediction errors can create an increasing pressure to actively shape one’s environment (or ‘niche’) to reduce or prevent these errors [36], following the idea of active (selective) evidence sampling, central to PP. As the avoidance generalizes, following the principles above, one’s world narrows, which in turn will be mirrored in a narrowing of cognition (one’s models). In a progressive, reciprocal fashion [37], this will again lead to a further shrinking of one’s world, and so on in a vicious mutually reinforcing feedback cycle. It is this gradual reciprocal narrowing of one’s goals and environments (cognition) that creates suffering and the need for psychotherapeutic support. Here, terms such as ‘disorder’ or ‘maladaptive behavior/beliefs’ will apply. Addiction, as well as anxiety and mood disorders can be usefully described in this fashion, where particular concerns come to dominate one’s world and collapse other goals and predictions. PP is uniquely suited to describe this developmental process, thereby reconceptualizing mental disorders as a possible, pernicious outcome of normal environmentally embedded learning, instead of as brain diseases [37]. This may cast a new light on the fact that no discrete biomarkers have been found (e.g. for affective disorders or addiction), and that many mental disorders share genetic risk factors [38,39] as well as a common psychological vulnerability factor [40]. It also helps us understand why pathogenetic reciprocal narrowing is more likely when living conditions have already started to narrow one’s paths available in the world (cf. the role of early adversity in life or of prior physical illness in mental disorder).

Toward more effective treatments

Current psychotherapeutic practice of affective disorders predominantly involves trying to uncover and modify a client’s maladaptive beliefs (CBT). The PP

account fits well in this tradition [41] because it provides a neurocognitive process theory about how these beliefs are formed and how they can be changed (using scaffolded learning environments). Correspondingly, most papers that cast psychopathology in PP terms emphasize belief updating [42] and, because this precision-based updating is thought to be mediated by neuromodulators, they express the hope for better pharmacological treatments in the future [43]. However, PP’s capacity to help us understand and improve psychotherapy is less explored so far. Here, we can only briefly touch on three important elements.

First, clinical practice should give attention to specifying the generative model of the client, to determine what the client construes as valuable (or ‘expected’) instead of relying on objective definitions or normative ideas of what is rewarding or unpleasurable (e.g. my idea [expectation] of what a mutually beneficial [valuable] social interaction entails might be very different from yours, which gives me significant anxiety when our pattern of interaction deviates from it). Failing to do so may make us either blind to the attractiveness of “what should objectively be” aversive situations (“Why does a person sometimes persist in ‘unrewarding’ behavior?” [44]), or puzzled about someone’s avoidance of “what should objectively be” neutral or even positive situations. For instance, harmless situations (e.g. asking a barista for a coffee) can, through the workings of our predictive mind, be construed as (near) catastrophes (e.g. the fact that she didn’t understand me the first time is proof that I am totally insignificant, which brings me physical anguish, which de facto makes it a dreadful experience instead of a harmless one). This of course precludes the updating of maladaptive avoidance behavior in exposure therapy [41].

Second, ‘model validation’ will be an important, recurrent step in psychotherapy. This means providing support (in dry PP terms: model evidence) for core constructs such as adaptive beliefs, values, and goals (cf. acceptance and commitment therapy). In fact, model validation is what we colloquially mean by ‘consolation’ (by a therapist or friend), that is, the reassurance that one’s emotional dynamics are to be expected (‘normal’) given the reference frame of the person (their beliefs, experienced events, and goals). It renders their experiences *predictable* and meaningful, rather than aberrant, irrational, or unpredictable. Earlier, we alluded to the ‘protective belt’ that models build for themselves by explaining away prediction errors using auxiliary hypotheses or by downregulating the salience of conflicting evidence by lowering our trust (expected confidence or certainty) in particular information sources. To alter these models, it is important to cultivate (renewed) epistemic trust in suppressed information sources and in one’s own actions, evoking a generalized sense of initiative beyond the therapeutic setting [45,46]. In PP

terms, this constitutes regaining confidence that one's actions will minimize prediction errors (that one's planning works, that a future can be produced). The model validation stage is crucial for regaining this trust/confidence.

Third, the model validation provides a safe basis for the introduction of new, more adaptive constructs and inferences in mental models. Here too, taking into account automatic predictions has shown to provide an advance beyond traditional association formation and behavioral change (CBT) ideas [47]. Eliciting a change in generative models requires us to set-up the environment to evoke prediction errors and to aid accommodation of these errors in an adaptive, sustainable manner. Simply put, therapists should create learning environments that allow clients to confidently predict that they will perform alternative behavior (e.g. drinking water in case of alcohol addiction) in relevant contexts (e.g. after dinner) and infer precise supporting beliefs for this behavior, most notably that this alternative behavior aligns better with their goals [48]. Together, the above three elements should help to widen (instead of narrowing) one's prospects again—the diversity of one's predictions, goals, and 'action radius' in the world.

Conclusion

The emerging description of mental distress in PP terms dovetails with insights from several psychotherapeutic traditions, as well as with dimensional (non-essentialist) and transdiagnostic views in psychopathology. The PP approach proposes plausible (inferential) mechanisms not only for pharmacological intervention, but also for the science of psychotherapy, showing that the 'soft' hermeneutic skills of psychotherapy [49] lend themselves to a 'dry' Bayesian PP analysis. It can do so because it is decidedly *not* biologically reductionist, through its emphasis on how the environment shapes inferences and constructed values [50]. Much more than a PP gloss on old ideas, the framework should continue to show great heuristic power for psychopathology.

Funding

This work is supported by the Flemish Government under Grant METH/14/02 and by the Scientific Research Foundation Flanders under Grant FWO19/PDS/041 awarded to Pieter van Dessel.

Conflict of interest statement

Nothing declared.

References

Papers of particular interest, published within the period of review, have been highlighted as:

- * of special interest
- ** of outstanding interest

1. Clark A: *Surfing uncertainty: prediction, action, and the embodied mind*. Oxford University Press; 2015. <https://market.android.com/details?id=book-ECgAAQBAJ>.
2. Hohwy J: *New directions in predictive processing*. Mind Lang; 2020. <https://onlinelibrary.wiley.com/doi/abs/10.1111/mila.12281>.
3. Friston K: **Embodied inference: or "I think therefore I am, if I am what I think,"**. In Tschacher W. *The implications of embodiment: cognition and communication*. Charlottesville, VA: Imprint Academic; 2011:89–125. <https://psycnet.apa.org/fulltext/2014-14659-005.pdf>.
4. Bruner JS: **On perceptual readiness**. *Psychol Rev* 1957, **64**:123. <http://psycnet.apa.org/journals/rev/64/2/123/>. Accessed 24 January 2013.
5. Kelly GA: **A brief introduction to personal construct theory**. *Perspectives in Personal Construct Theory* 1970, **1**:29. <https://www.aipcc.it/wp-content/uploads/2019/04/2017.01.003.025.pdf>.
6. von Glasersfeld E. *Radical constructivism: a way of knowing and learning*. *Studies in mathematics education series*, vol. 6. Bristol, PA: Falmer Press, Taylor & Francis Inc.; 1995. <http://eric.ed.gov/?id=ED381352>. Accessed 25 September 2014.
7. Hoffman DD, Singh M, Prakash C: **The interface theory of perception**. *Psychon Bull Rev* 2015, **22**:1480–1506, <https://doi.org/10.3758/s13423-015-0890-8>.
8. Smith R, Friston K, Whyte C: *A step-by-step tutorial on active inference and its application to empirical data*. 2021, <https://doi.org/10.31234/osf.io/b4j6>.
9. Gershman SJ: **What does the free energy principle tell us about the brain?** *arXiv [q-bio.NC]* 2019. <http://arxiv.org/abs/1901.07945>; 2019.
10. Bogacz R: **A tutorial on the free-energy framework for modeling perception and learning**. *J Math Psychol* 2015. <http://www.sciencedirect.com/science/article/pii/S0022249615000759>; 2015.
11. Hohwy J: *The self-evidencing brain*. Noûs; 2014, <https://doi.org/10.1111/nous.12062>.
12. Bem DJ: **Self-perception theory**. *Adv Exp Soc Psychol* 1972, **6**: 1–62. <http://healthyinfluence.com/wordpress/wp-content/uploads/2011/05/SP-Theory-Bem-Advances.pdf>.
13. Pulcu E, Browning M: **The misestimation of uncertainty in affective disorders**. *Trends Cognit Sci* 2019, **23**:865–875, <https://doi.org/10.1016/j.tics.2019.07.007>.
- ** The authors review evidence that estimation of uncertainty plays an important role in learning and explain how misestimation of uncertainty can lead to symptoms of anxiety and depression and might underlie affective disorders.
14. Moors A, Boddez Y, De Houwer J: **The power of goal-directed processes in the causation of emotional and other actions**. *Emot Rev* 2017, **9**:310–318, <https://doi.org/10.1177/1754073916669595>.
15. Van Dessel P, Hughes S, De Houwer J: **How do actions influence attitudes? An inferential account of the impact of action performance on stimulus evaluation**. *Pers Soc Psychol Rev* 2019, **23**:267–284, <https://doi.org/10.1177/1088868318795730>.
16. Austin JT, Vancouver JB: **Goal constructs in psychology: structure, process, and content**. *Psychol Bull* 1996, **120**:338. <https://psycnet.apa.org/doiLanding?doi=10.1037/0033-2909.120.3.338>.
17. Pezzulo G, Rigoli F, Friston K: **Active Inference, homeostatic regulation and adaptive behavioural control**. *Prog Neurobiol* 2015, **134**:17–35, <https://doi.org/10.1016/j.pneurobio.2015.09.001>.
18. FitzGerald THB, Dolan RJ, Friston KJ: **Model averaging, optimal inference, and habit formation**. *Front Hum Neurosci* 2014, **8**: 457, <https://doi.org/10.3389/fnhum.2014.00457>.
19. Clark A: **Beyond desire? Agency, choice, and the predictive mind, australas**. *J Philos* 2020, **98**:1–15, <https://doi.org/10.1080/00048402.2019.1602661>.
20. Kommers C: *The intuitive anthropologist: why "intuitive psychology" falls short for making sense of those who are different*. 2020, <https://doi.org/10.31234/osf.io/3sc96>.

21. Van de Cruys S, Evers K, Van der Hallen R, Van Eylen L, Boets B, de-Wit L, Wagemans J: **Precise minds in uncertain worlds: predictive coding in autism.** *Psychol Rev* 2014, **121**: 649–675, <https://doi.org/10.1037/a0037665>.
22. Sterzer P, Adams RA, Fletcher P, Frith C, Lawrie SM, Muckli L, Petrovic P, Uhlhaas P, Voss M, Corlett PR: **The predictive coding account of psychosis.** *Biol Psychiatr* 2018, **84**:634–643, <https://doi.org/10.1016/j.biopsych.2018.05.015>.
23. Fradkin I, Adams RA, Parr T, Roiser JP, Huppert JD: **Searching for an anchor in an unpredictable world: a computational model of obsessive compulsive disorder.** *Psychol Rev* 2020, **127**:672–699, <https://doi.org/10.1037/rev0000188>.
24. Duncan S, Barrett LF: **Affect is a form of cognition: a neurobiological analysis.** *Cognit Emot* 2007, **21**:1184–1211, <https://doi.org/10.1080/02699930701437931>.
25. Frijda NH: **The laws of emotion.** *Am Psychol* 1988, **43**:349–358, <https://doi.org/10.1037/0003-066X.43.5.349>.
26. Hesp C, Smith R, Parr T, Allen M, Friston KJ, Ramstead MJD: **Deeply felt affect: the emergence of valence in deep active inference.** *Neural Comput* 2021, **33**:398–446, https://doi.org/10.1162/neco_a_01341.
27. Van de Cruys S: **Affective value in the predictive mind.** In *Philosophy and predictive processing*. Edited by Metzinger TK, Wiese W, Frankfurt am Main: MIND Group; 2017, <https://doi.org/10.15502/9783958573253>.
28. Joffily M, Coricelli G: **Emotional valence and the free-energy principle.** *PLoS Comput Biol* 2013, **9**, e1003094, <https://doi.org/10.1371/journal.pcbi.1003094>.
29. Whalen PJ: **The uncertainty of it all.** *Trends Cognit Sci* 2007, **11**: 499–500, <https://doi.org/10.1016/j.tics.2007.08.016>.
30. Muth C, Raab MH, Carbon C-C: **The stream of experience when watching artistic movies. Dynamic aesthetic effects revealed by the Continuous Evaluation Procedure (CEP).** *Front Psychol* 2015, **6**:365, <https://doi.org/10.3389/fpsyg.2015.00365>.
31. Huys QJM, Dayan P: **A Bayesian formulation of behavioral control.** *Cognition* 2009, **113**:314–328, <https://doi.org/10.1016/j.cognition.2009.01.008>.
32. Ellis BJ, Bianchi J, Griskevicius V, Frankenhuis WE: **Beyond risk and protective factors: an adaptation-based approach to resilience.** *Perspect Psychol Sci* 2017, **12**:561–587, <https://doi.org/10.1177/1745691617693054>.
33. Taylor SE, Brown JD: **Illusion and well-being: a social psychological perspective on mental health.** *Psychol Bull* 1988, **103**:193–210. <https://www.ncbi.nlm.nih.gov/pubmed/3283814>.
34. Kwisthout J, Bekkering H, van Rooij I: **To be precise, the details don't matter: on predictive processing, precision, and level of detail of predictions.** *Brain Cognit* 2017, **112**:84–91, <https://doi.org/10.1016/j.bandc.2016.02.008>.
35. Van den Bergh O, Brosschot J, Critchley H, Thayer JF, Ottaviani C: **Better safe than sorry: a common signature of general vulnerability for psychopathology.** *Perspect Psychol Sci* 2020, <https://doi.org/10.1177/1745691620950690>. 1745691620950690.
The authors explain how a person's general strategy to process prediction error in low detail can give rise to weakly informed prior beliefs that might underlie (affective) mental disorders.
36. Constant A, Bervoets J, Hens K, Van de Cruys S: **Precise worlds for certain minds: an ecological perspective on the relational self in autism.** *Topoi* 2018, <https://doi.org/10.1007/s11245-018-9546-4>.
37. Lewis M: **Brain change in addiction as learning, not disease.** *N Engl J Med* 2018, **379**:1551–1560, <https://doi.org/10.1056/NEJMra1602872>.
- The author reviews neurobiological alterations in addiction and concludes they cannot be labelled as pathological, let alone specific to the disorder. He presents an alternative learning-based account of addiction based on the concept of reciprocal narrowing that may fruitfully be applied to mental disorders more generally.
38. Gratten J, Wray NR, Keller MC, Visscher PM: **Large-scale genomics unveils the genetic architecture of psychiatric disorders.** *Nat Neurosci* 2014, **17**:782–790, <https://doi.org/10.1038/nn.3708>.
39. Cross-Disorder Group of the Psychiatric Genomics Consortium: **Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis.** *Lancet* 2013, **381**:1371–1379, [https://doi.org/10.1016/S0140-6736\(12\)62129-1](https://doi.org/10.1016/S0140-6736(12)62129-1).
40. Caspi A, Moffitt TE: **All for one and one for all: mental disorders in one dimension.** *Am J Psychiatr* 2018, **175**:831–844, <https://doi.org/10.1176/appi.ajp.2018.17121383>.
41. Moutoussis M, Shahar N, Hauser TU, Dolan RJ: **Computation in psychotherapy, or how computational psychiatry can aid learning-based psychological therapies.** *Comput Psychiatr* 2018, **2**:50–73, https://doi.org/10.1162/CPSY_a_00014.
The authors describe how recent advances in PP and computational modeling can help formalize and test hypotheses regarding the maladaptive inferences that underlie mental disorder and can improve learning-based therapies such as CBT.
42. Kube T, Rozenkrantz L: **When beliefs face reality: an integrative review of belief updating in mental health and illness.** *Perspect Psychol Sci* 2020, <https://doi.org/10.1177/1745691620931496>. 1745691620931496.
The authors review evidence about how healthy people and people with mental disorders update their beliefs after receiving new information that supports or challenges their views. They argue that the strategy to afford very high or very low precision to prior beliefs may underlie psychopathology.
43. Constant A, Hesp C, Davey CG, Friston KJ, Badcock PB: **Why depressed mood is adaptive: a numerical proof of principle for an evolutionary systems theory of depression.** *Comput Psychiatr* 2021, **5**:60–80, <https://doi.org/10.5334/cpsy.70>.
44. Kelly GA: *Man's construction of his alternatives. Assessment of Human Motives*; 1958:33–64.
45. Luyten P, Campbell C, Fonagy P: **Rethinking the relationship between attachment and personality disorder.** *Curr Opin Psychol* 2021, **37**:109–113, <https://doi.org/10.1016/j.copsyc.2020.11.003>.
46. Hunt JM: **Experiential roots of intention, initiative, and trust.** In *Advances in intrinsic motivation and aesthetics*. Edited by Day HI, Boston, MA: Springer US; 1981:169–202, https://doi.org/10.1007/978-1-4613-3195-7_8.
47. Van Dessel P, Hughes S, De Houwer J: **Consequence-based approach-avoidance training: a new and improved method for changing behavior.** *Psychol Sci* 2018, **29**:1899–1910, <https://doi.org/10.1177/0956797618796478>.
48. Wiers RW, Van Dessel P, Köpetz C: **ABC training: a new theory-based form of cognitive-bias modification to foster automatization of alternative choices in the treatment of addiction and related disorders.** *Curr Dir Psychol Sci* 2020, **29**: 499–505. <https://journals.sagepub.com/doi/abs/10.1177/0963721420949500>.
49. Chessick RD: **Hermeneutics for psychotherapists.** *Am J Psychother* 1990, **44**:256–273, <https://doi.org/10.1176/appi.psychotherapy.1990.44.2.256>.
50. Adams RA, Huys QJM, Roiser JP: **Computational Psychiatry: towards a mathematically informed understanding of mental illness.** *J Neurol Neurosurg Psychiatry* 2016, **87**:53–63, <https://doi.org/10.1136/jnnp-2015-310737>.